

1 **The measurement of skin thickness in Merino sheep using real time ultrasound**

2 *D.J. Brown^{1,2}, M.L. Wolcott³ and B.J. Crook¹*

3 ¹ Animal Science, University of New England, Armidale, NSW, 2351

4 ² Present Address: Animal Genetics and Breeding Unit, University of New England, Armidale, NSW, 2351

5 ³ Cattle and Beef Industry CRC, University of New England, Armidale, NSW, 2351

6

7 Short Title: Measurement of skin thickness in sheep

8

9 Contact address:

10 Daniel Brown

11 Animal Genetics and Breeding Unit,

12 University of New England,

13 Armidale NSW 2351 Australia

14 Ph: (02) 6773 2160

15 Fax: (02) 6773 3266

16 email: dbrown2@metz.une.edu.au

17

18

19

1 **Summary**

2
3 The measurement of skin thickness in sheep may offer a number of potential benefits for sheep
4 producers in terms of wool growth and monitoring body condition. The relationship between skin
5 thickness measurement made using skin fold callipers to those made using real time ultrasound
6 equipment was compared in 16 grazing Merino wethers. The relationship between the
7 measurements made using the two techniques varied throughout the year. The measurements were
8 not significantly different ($P < 0.05$) at 7 of the 12 measurement times throughout the year. The
9 measurement techniques were also moderately to highly correlated at 6 of the 12 measurement
10 times however these were not exactly the same times as those with means not significantly
11 different. With modified equipment, accurate estimation of average skin thickness could be
12 measured using ultrasound.

13 **Keywords:** Skin fold callipers, real time ultrasound, seasonal variation

14 15 **Introduction**

16
17 It has been suggested that differences between sheep in skin thickness may be associated with
18 differences in wool production. Utilising skin thickness may improve genetic selection of wool
19 growth rate and quality. Skin thickness is positively related to clean fleece weight (0.39 ± 0.13),
20 mean fibre diameter (0.20 ± 0.11) and staple length (0.35 ± 0.12) (Gregory 1982b). These
21 relationships have been reviewed in more detail by Hynd *et al.* (1997). Sheep selected for high
22 fleece weight also have a thicker skin than those selected for reduced fleece weight (Williams and
23 Thornberry 1992). Skin thickness was not related to the differences in liveweight between the two
24 groups. When individual animals from these groups were pen fed, these differences were not
25 observed (Williams and Thornberry 1992) suggesting that diet and or climatic conditions may
26 influence these relationships. Significant differences in skin thickness have also been observed

1 between bloodlines of Merino sheep (Murray 1996). In contrast to these results, Masters *et al.*
2 (2000) found no significant differences between sheep of similar mean fibre diameter but selected
3 for and against fleece weight. As skin thickness is a heritable trait (0.25 to 0.60) (Gregory 1982a;
4 Slee *et al.* 1991) it is anticipated that this trait would respond to selection.

5
6 It has been shown that the weight of the skin per unit area shows similar seasonal trends to that of
7 body weight (Hutchinson 1957; Wodzicka 1958b; Lyne 1964; Williams and Morley 1994; Murray
8 1996; Schlink *et al.* 1996; Briegel *et al.* 2000). It is possible that differences in average skin
9 thickness and seasonal variation in skin thickness between animals is associated with differences in
10 the environmental responsiveness of wool growth. Average skin thickness and seasonal variation in
11 skin thickness are significantly correlated with characteristics of the fibre diameter profile, mean
12 fibre diameter, variation in fibre diameter, staple length and staple strength (Brown and Crook
13 2000). Sheep that are able to maintain skin thickness in spite of decreases in body condition may be
14 able to maintain superior follicle nutrition and wool growth.

15
16 While the most widely used and accepted technique for measuring skin thickness is to measure the
17 thickness of a fold of skin with skin-fold callipers, variation in the levels of subcutaneous fat and
18 pinching force can contribute to measurement error (Alexander and Miller 1979). Ultrasound
19 scanning is an objective method for estimating subcutaneous fat thickness and is frequently used in
20 both sheep and cattle (Dicker *et al.* 1988). Ultrasound scanning provides an objective, non-invasive
21 and simple way to fat and muscle depths and potentially skin thickness. Ultrasound also provides a
22 real-time image of the skin, its structure and characteristics. The high frequency, high-resolution
23 ultrasonic echo technique is capable of determining human skin thickness quickly and accurately
24 (Alexander and Miller 1979).

1 Butler (1991) noted that variation may exist between sheep in the depth and relative proportions of
2 epidermis, reticular layer and papillary layer of the dermis, sebaceous gland, sudoriferous gland and
3 echo-pattern. These differences may influence wool production and the environmental
4 responsiveness and fibre diameter profiles of sheep. By using high frequency and high-resolution
5 ultrasound scanning differences between sheep in these characteristics may be identified.

6
7 This study was designed to examine the relationship between skin thickness measurements made
8 using skin fold callipers and those using real time ultrasound equipment throughout the year.

9 10 **Materials and Methods**

11
12 A mob of 16 2-year-old fine wool Merino wethers were maintained for 12 months on improved
13 pastures at Armidale NSW. Skin thickness was measured using both skin fold callipers and
14 ultrasound at 12 times throughout the experiment, approximately 4 weeks apart.

15
16 The calliper technique was a modification of the technique used by Williams and Thornberry
17 (1992). Skin thickness was measured at two points randomly selected on the right mid-side after
18 close clipping (Oster small animal clippers size 30 blade). These two points were marked with a
19 permanent-marking pen and used for the skin measurements at each sampling. A double fold of
20 skin was measured using dial-gauge callipers that exerted a constant pressure of 1250g/cm² (Lyne
21 1964). A measurement was made at each site with the callipers facing in anterior / posterior
22 direction relative to the body of the sheep and a second measurement was made in a dorsal / ventral
23 direction. Average skin thickness at each sampling time was calculated by dividing the sum of
24 these measurements by 8 (2 measurements x 2 locations x 2 layers).

25

1 The thickness of skin was also measured with a real time ultrasound scanner at the C site (over the
2 eye muscle at the 12th rib) at the same time as the calliper measurements. These measurements
3 were made using an Aloka 500V real time ultrasound scanner at a frequency of 3.5 Mega Hertz.
4 The probe was 17.5cm long and designed for use on cattle. The existing wool staples at the C site
5 were separated to gain access to the skin and vegetable oil was applied to the skin to enable clear
6 image transmission. The site was not clipped as clipping has been shown to influence skin thickness
7 (Wodzicka 1958c; Lyne 1964).

8
9 For each animal mean skin thickness and variation (standard deviation and coefficient of variation)
10 in skin thickness over the experimental period were calculated for both techniques.

11
12 To compare measurements, least squares multivariate analysis of variance of the skin thickness
13 measurements was conducted using the General Linear Model procedure in SAS (1990). Residual
14 partial correlation coefficients were calculated to compare the skin thickness measurements from
15 the calliper and ultrasound techniques.


16 17 **Results**

18
19 The means, standard errors and correlation coefficients for each skin thickness measurement
20 technique at each sampling time are shown in Table 1. The relationship between the calliper and
21 ultrasound techniques for estimating skin thickness varied throughout the experiment (Figure 1).

22

23

1 **Figure 1 The mean values for the ultrasound (▲) and skin fold calliper (■) techniques**
2 **throughout the experiment**



3
4 The ultrasound measurements were significantly ($P<0.05$) higher than the calliper measurements in
5 the months of December, January, April and May while significantly ($P<0.05$) lower in June. The
6 measurements made in the remaining months were not significantly ($P>0.05$) different from the
7 measurements made using the calliper technique. The two techniques were highly correlated in
8 July ($r=0.84$) and moderately correlated in February, April, June, September, October and
9 November ($r=0.41$ to 0.66). The mean, standard deviation and coefficient of variation of the
10 measurements throughout the experiment were significantly different ($P<0.05$) between the calliper
11 and ultrasound techniques but moderately and significantly correlated ($r=0.66$, 0.51 to 0.56
12 respectively). Both Figure 1 and Table 1 suggest that the accuracy of the ultrasound technique in
13 measurement skin thickness improved throughout the experiment.

14

15

Table 1 The mean values (mm), standard errors of the means (s.e.) and correlation coefficients for the relationship between the skin thickness measurement made using the calliper and those made using real time ultrasound technique throughout the experiment


Month	Callipers	Ultrasound	s.e.	Correlation	Month	Callipers	Ultrasound	s.e.	Correlation
Dec	1.77 ^a	2.25 ^b	0.11	0.05	Aug	1.73 ^a	1.56 ^a	0.08	0.19
Jan	1.74 ^a	2.56 ^b	0.13	0.09	Sep	1.83 ^a	1.75 ^a	0.07	0.41
Feb	1.85 ^a	1.91 ^a	0.06	0.46*	Oct	1.73 ^a	1.81 ^a	0.06	0.55*
Mar	1.82 ^a	2.00 ^a	0.10	0.07	Nov	1.77 ^a	1.84 ^a	0.07	0.50*
Apr	1.72 ^a	2.16 ^b	0.06	0.18	Mean	1.77 ^a	1.93 ^b	0.03	0.66*
May	1.78 ^a	2.06 ^b	0.06	0.24	SD	0.11 ^a	0.47 ^b	0.03	0.51*
Jun	1.76 ^a	1.50 ^b	0.04	0.66*	CV	6.15 ^a	24.22 ^b	1.14	0.56*
Jul	1.76 ^a	1.75 ^a	0.03	0.84*					

Means within each column with different superscripts are significantly different (P<0.05)

* Correlation significant (P<0.05)

Figure 2 illustrates the relationship between the mean skin thickness measurements from the ultrasound technique and those from the calliper technique. This graph suggests that the positive bias of the ultrasound technique is more prominent in sheep with thinner skins.

Figure 2 The relationship between mean skin thickness measured using callipers and mean skin thickness measured using real time ultrasound



1 **Discussion**

2

3 The relationship between calliper and ultrasound measurements of skin thickness varied throughout
4 the year and were only highly correlated at the eighth sampling time. However five of the monthly
5 measurements were significantly correlated ($r=0.46$ to 0.84) suggesting that the ultrasound
6 measurements were similar to those made by the callipers. A number of factors may have
7 influenced these results. The ultrasound equipment used in this experiment was designed for the
8 measurement of fat and eye muscle depth in cattle. The machine operated at a relatively low
9 frequency and is designed to measure larger ranges. As the skin of sheep is very thin, generally less
10 than 2mm, the image on this machine was very small. This made identification of the various tissue
11 layers difficult. Furthermore the ultrasound equipment only measured to the nearest 0.5mm which
12 equates to approximately 25% of the overall thickness of the skin. The skin fold callipers measured
13 to the nearest 0.1mm, which resulted in a greater ability to distinguish differences between sheep.
14 Subcutaneous fat measurements can be made using conventional scanners operating at 5 to 7 MHz
15 (Butler 1991). Ultrasound equipment operating at higher frequency and resolution are required to
16 produce adequate images to identify features of the skin (Alexander and Miller 1979; Dines *et al.*
17 1984). As the equipment used in this study operated at 3.5 MHz it is anticipated that these
18 differences influenced the relationship between the ultrasound and calliper based measurements.

19

20 Results suggest that the accuracy of the ultrasound method improved as the experiment progressed.
21 This may have been a direct result of the experience gained from the measurements at the previous
22 sampling times. As the experiment progressed the operator found it easier to identify the layers of
23 the dermis, subcutaneous fat and muscle and therefore to judge the top and bottom surfaces of the
24 skin. Further experimentation may benefit from the incorporation of a leading in period to adapt
25 operators to measurement and perhaps calibration against calliper measurements.

26

1 The two techniques used a different site to measure skin thickness. It is well known that wool traits
2 vary over the body of the sheep (Sumner and Craven 2000). Variations in skin thickness of the
3 body of sheep are not well documented, however, significant differences in skin thickness have
4 been observed between sites in Merino sheep (Wodzicka 1958a; Williams and Thornberry 1992).
5 Associated with this, to enable accurate measurements the site at which the calliper measurements
6 were made had to be clipped prior to measurement whereas the ultrasound measurement did not
7 always require the site to be clipped, except of rare occasions when the oil reacted unfavorably with
8 the wool. As clipping has been shown to influence skin thickness (Wodzicka 1958c; Lyne 1964)
9 this may have influenced the relationship between the measurement from the two techniques.

10
11 The fact that the wool generally does not have to be clipped is another benefit of the ultrasound
12 technique. Clear images could be obtained simply by parting the wool and applying a small amount
13 of vegetable oil. In a small percentage of animals (12.5%) however, the oil reacted unfavorably
14 with the wool, discoloring it and creating a potential fly-strike problem. The above average rainfall
15 that was received throughout the experiment may have influenced these reactions. It may be that the
16 use of water based scanning gels, available for medical applications, would provide a more
17 appropriate couplant for use in sheep.

18 19 **Conclusion**

20
21 This study found that real time ultrasound scanning could measure skin thickness and offered many
22 potential advantages over the calliper technique. Further research is required to identify the most
23 appropriate real time ultrasound equipment for this application, and it is likely to produce a viable
24 alternative to the current skin fold calliper technique.

1 **Acknowledgments**

2
3 The authors would like to thank the CSIRO Division of Animal Production at Prospect for
4 providing the skin fold callipers for this research. The authors would also like to acknowledge Mr.
5 Chris Shands for his assistance with ultrasound scanning. D.Brown was in receipt of an Australian
6 government scholarship and was also partly funded by the Cooperative Research Center for
7 Premium Quality Wool.

9 **References**

- 10
11 Alexander, H., and Miller, D. L. (1979). Determining skin thickness with pulsed ultra sound,
12 *Journal of Investigative Dermatology* **72(1)**, 17-19.
- 13 Briegel, J.R., Liu, S.M. and Adams, N.R. (2000) Skin protein mass and protein synthesis rate
14 contributes equally to seasonal increase in wool growth, *The Asian-Australian Journal*
15 *of Animal Science* **23**, 153.
- 16 Brown, D.J. and Crook, B.J. (2000). Determinants of environmental responsiveness of fibre
17 diameter in grazing fine wool Merino sheep, *Australian Journal of Agricultural*
18 *Research* (Submitted).
- 19 Butler, L. G. (1991). Potential application of new technologies in research and as indirect selection
20 criteria for wool production efficiency or quality, In "Wool Biology", Ed. P. I. Hynd,
21 page, 106-110, Australian Wool Corporation.
- 22 Cottle, D. J. (1991). "The Australian Sheep and Wool Handbook" Inkata Press : Sydney.
- 23 Dicker, R. W., Fowler, D. G., Perry, D., and Sundstrom, B. (1988). Accuracy of real-time
24 ultrasound systems for fat thickness estimation in live cattle, *Proceedings of the*
25 *Australian Society of Animal Production* **17**, 178-181.

1 Dines, K. A., Sheets, P. W., Brink, J. A., Hanke, C. W., Condra, K. A., Clendenon, J. L., Goss, S.
2 A., Smith, D. J., and Franklin, T. D. (1984). High frequency ultrasonic imaging of skin:
3 Experimental results, *Ultrasonic Imaging* **6**, 408-434.

4 Gregory, I. P. (1982a). Genetic Studies of South Australian Merino Sheep III Heritabilities of
5 Various Wool and Body Traits, *Australian Journal of Agricultural Research* **33**, 355-
6 362.

7 Gregory, I. P. (1982b). Genetic Studies of South Australian Merino Sheep IV Genetic, Phenotypic
8 and Environmental Correlations Between Various Wool and Body Traits, *Australian*
9 *Journal of Agricultural Research* **33**, 363-373.

10 Hutchinson, K. J. (1957). Changes in the thickness, weight and protein content of the skin of two
11 adult Merino ewes, *Journal of the Australian Institute of Agricultural Science* **23**, 238-
12 241.

13 Hynd, P.I., Ponzoni, R.W. and Hill, J.A. Hill (1997) Can selection for skin traits increase the rate of
14 genetic progress in Merino breeding programs, *Association for the Advancement of*
15 *Animal Breeding and Genetic*, **12(2)**, 752-759.

16 Lyne, A. G. (1964). Effect of adverse nutrition on the skin and wool follicles in Merino sheep,
17 *Australian Journal of Agricultural Research* **15**, 788-801.

18 Masters, D.G., Liu, S.M., Purvis, I.W. and Hartofillis, M. (2000) Wool growth and protein synthesis
19 in the skin of superfine Merinos with high and low fleece-weight, *The Asian-Australian*
20 *Journal of Animal Science* **23**, 457-460.

21 Murray, P.J. (1996). The effect of nutrition on skin thickness of fine and broad wool Merino sheep,
22 *Proceedings of the Australian Society of Animal Production* **21**, 388.

23 SAS (1990). SAS/STAT User's Guide. Version 6, Fourth edition, Cary, N.C. : SAS Institute.

24 Schlink, T., Adams, N., and Peterson, A. (1996). Components of staple strength, In "*Proceedings of*
25 *the Cooperative Research Center Roadshow*", page 29, Cooperative Research Center
26 for Premium Quality Wool.

- 1 Slee, J., Alexander, G., Bradley, L.R., Jackson, N. and Stevens, D. (1991) Genetic aspects of cold
2 resistance and related characters in newborn Merino lambs, *Australian Journal of*
3 *Experimental Agriculture* **31(2)**, 175-182.
- 4 Sumner, R.M.W. and Craven, A.J. (2000). Variation of fibre and follicle characteristics related to
5 wool bulk over the body of Perendale ewes: Implications for measurement of wool bulk,
6 *Proceedings of the New Zealand Society of Animal Production* **60**, 166-170.
- 7 Williams, A. J., and Thornberry, K. J. (1992). The skin thickness of medium wool Merino sheep
8 and its relationship to wool production, *Proceedings of the Australian Society of Animal*
9 *Production* **19**, 138-141.
- 10 Williams, A. J. and Morley, F.C. (1994). Influence of dietary intake and genetic capacity for wool
11 growth on the composition of mid-trunk skin of Merino sheep, *Australian Journal of*
12 *Agricultural Research* **45(8)**, 1715-1729.
- 13 Wodzicka, M. (1958a). Studies on the thickness and chemical composition of the skin of sheep I –
14 Development techniques, *New Zealand Journal of Agricultural Research* **1**, 582-591.
- 15 Wodzicka, M. (1958b). Studies on the thickness and chemical composition of the skin of sheep II –
16 Variations during growth, *New Zealand Journal of Agricultural Research* **1**, 592-600.
- 17 Wodzicka, M. (1958c). Studies on the thickness and chemical composition of the skin of sheep III –
18 Effect of shearing, *New Zealand Journal of Agricultural Research* **1**, 601-606.
- 19